

CAUTION

KEEP OUT OF REACH OF CHILDREN


USE WITHIN 24 MONTHS FROM
DATE OF MANUFACTURE



CoRoN[®]

Foliar Fertiliser 30-0-0 (w/v) Nitrogen (N) - 30% w/v

A Liquid Controlled Release
Nitrogen Fertiliser



DISTRIBUTED BY:
Campbells Fertilisers Australasia Pty Ltd
18 Raymond Road, Laverton North,
Victoria 3026 Australia
Phone: +61 3 9931 2211 Fax: +61 3 9931 2201
www.cambellsfert.com.au



AUTHORISED DISTRIBUTOR

Is a registered trademark of @CoRoN
is a registered trademark of Helena
Holding Company

PRECAUTION: Use of the product in accordance with label directions is not expected to adversely affect normal vegetative and/or reproductive growth in most situations. However, various environmental and agronomic factors, such as crop stress associated with high temperatures and/or dry conditions, make it impossible to eliminate all risks associated with the use of the product; therefore, crop injury is always possible. The user assumes all risks of use and handling. In an effort to reduce the risk of crop injury, apply early morning or late evening. Use in accordance with recommendations of a qualified individual or institution, such as, but not limited to, a certified crop advisor, agronomist, university crop extension publication, or apply according to recommendations in your approved nutrient management plan. Consult your local Campbells Fertilisers Australasia representative for further information.

GENERAL USE RATE RECOMMENDATIONS

FIELD CROPS

LUCERNE: 10-15L per hectare. Apply after each cutting when sufficient foliage is present.

CANOLA: 10-20L per hectare. Apply pre-bloom.

CORN (Field, Pop, Seed, Sweet): 30-50L per hectare at V-5 to V-8 as a foliar application for use as a partial side-dress Nitrogen (N) replacement. 10-30L per hectare applied as a foliar application at pre-tassel as a Nitrogen (N) supplement. 10-20L per hectare applied as a foliar application when used with fungicides, insecticides. Maximum replacement value of side-dress Nitrogen (N) using CoRoN® 30-0-0: 30-50L per hectare.

COTTON (SEEDLING): 8-10L per hectare. Apply when first true leaves appear. (AFTER SEEDLING STAGE): 5-10 L per hectare. (BOLL DEVELOPMENT): 10 -30 L per hectare. Apply at early boll formation and repeat at 14-21 day intervals.

GRAIN SORGHUM: 10-15 L per hectare after pollination.

PEANUTS: (Single Application) – 25-40L per hectare at early pod development. (Multiple Application) – 10-15L per hectare per application at early bloom and repeat at 14-21 day intervals until pods are filled.

POTATOES: (Single Application) – 25-40L per hectare. Apply at mid tuber development. (Multiple Application) – 10-15L per hectare at tuber initiation and repeat at 10-14 day intervals until maximum tuber development has occurred.

RICE: 15-25L per hectare. Apply at panicle emergence.

WHEAT, BARLEY, OATS: 10-30L per hectare from tillering to flag leaf stage as a Nitrogen (N) supplement. Maximum replacement value for partial Nitrogen (N) replacement: 30L per hectare. Nitrogen (N). Certain conditions may change the Maximum replacement value, but only with the approval of your Campbells sales representative using prior experience and local data.

SOYBEANS: 10-30L per hectare as a Nitrogen (N) supplement. Apply at V6-R4 growth stage.

SUGARCANE: 20-40L per hectare. Apply from tillering to out of hand growth stage.

SUNFLOWER: 10-15L per hectare. Apply when outer seeds start to fill, repeat in 10-14 days.

SWEET POTATOES: 10-15L per hectare. Apply at tuber initiation and repeat at 10-14 days.

OTHER CROPS: 10-15L per hectare. Apply when sufficient foliage is present, try on a small area until more experience and trials have been completed to determine if higher rate is desirable.

VEGETABLES

ASPARAGUS: 15-25L per hectare. Apply beginning at mid-fern development, then at 14-21 day intervals.

BEANS (DRY): (Single Application) – 25L per hectare at early pod formation. (Multiple Application) – 10-15L per hectare at early full flower and repeat in 10-14 days.

BEANS (GREEN, LIMA): 10-15L per hectare at early flower and repeat in 7-10 days.

BROCCOLI: 15-25L per hectare. Apply first application after thinning, then 3 weeks before head formation and repeat in 7-10 days.

CABBAGE: 15-25L per hectare Apply first application after thinning, then early head formation and repeat after 14-21 days. **MELONS:** 15-25L per hectare. Apply at early flowering and repeat 7-10 days later.

CAULIFLOWER: 15-25L per hectare. Apply first application after thinning or transplant, other applications at early head set and repeat at 10-14 day intervals.

CELERY: 10-15L per hectare. Apply when plants are 20-30 centimetres tall and repeat at 10-14 day intervals.

CUCUMBERS: (Single Application) – 25-40L per hectare. Apply at early fruit set. (Multiple Applications) – 10-15L per hectare. Apply at early flowering and repeat at 10-14 day intervals.

LENTILS: 10-15L per hectare. Apply at early flowering and repeat at 10-14 day intervals.
LETTUCE: 10-15L per hectare. Apply after thinning, then at early head formation and repeat at 10-14 day intervals.
ONIONS: 10-15L per hectare. Apply at mid set development and repeat at 14-21 day intervals.
PEAS: 10-15L per hectare. Apply at early flowering and repeat in 10-14 days.
CAPSICUM: 10-15L per hectare Apply at early fruit set and repeat in 10-14 days.
SPINACH: 15-25L per hectare. Apply when sufficient foliage is present and repeat at 14-21 days.
SQUASH: 15-25L per hectare. Apply at early fruit set and repeat at 10-14 day intervals.
TOMATOES (PROCESS): (Single Application) – 25L per hectare. Apply 10-14 days after full bloom. (Multiple Applications) – 10-15L per hectare at full bloom and repeat at 10-14 day intervals until harvest is near completion.
WATERMELONS: 15-25L per hectare. Apply at early flowering and repeat 7-10 days later.
OTHER CROPS: 10-15L per hectare. Apply when sufficient foliage is present, try on a small area until more experience and trials have been completed to determine if higher rate is desirable.

FRUITS AND NUTS

ALMONDS: Apply 15-25L per hectare at full leaf. Repeat at early nut expansion.
APPLES: Apply 10-15L per hectare prior to fruit set or post-harvest.
CHERRIES: Apply 10-15L per hectare prior to fruit set.
CITRUS: Apply 10-15L per hectare at early bloom and repeat after fruit set.
GRAPES: Apply 5-10L per hectare prior to fruit set. Application of nitrogen during fruit set can increase the risk of shatter.
PEACHES/NECTARINES: Apply 15-25L per hectare prior to fruit set.
PEARS: 10-15L per hectare prior to fruit set or post-harvest.
PECANS: Apply 10-15L per hectare at full leaf. Repeat at early nut expansion.
PLUMS: Apply 15-25L per hectare prior to fruit set.
OLIVES: Apply 10-15L per hectare at early fruit development and repeat as required.
STRAWBERRIES: Apply 5-8L per hectare at early flowering and repeat every 14 days through harvest.
OTHER CROPS: 10-15L per hectare. Apply when sufficient foliage is present, try on a small area until more experience and trials have been completed to determine if higher rate is desirable.

Mixing and Application:

Add ½ total water to spray tank. Turn on recirculation line. Add any pesticides and other materials. Add desired amount of CoRoN® 30-0-0. Bring water to final volume. Apply in adequate spray volume for complete coverage. For fertigation treatments, add concentrate to injection tank

Do not apply during extremes of climate, during intense heat or just prior to or during rain events.

Tank mixing or co-application:

Read all labels carefully and adhere strictly to instructions on mixing. Apply to small test area only when unfamiliar with products or new crops. Many variables are outside the control of the manufacturer and can influence results of co-applied products and therefore co-application is entirely at the risk of end user.

STORAGE:

Keep in cool, dry chemical store, out of the reach of children and animals. Protect from frost and other climatic extremes (including temperatures exceeding 40 • C). Poor water quality may reduce the effectiveness of the product, the solubility of the product and/or increase the potential for phytotoxic effects on plants

SAFETY DIRECTIONS:

Keep out of reach of children. Avoid release to the environment. Wear protective gloves/protective clothing/eye protection/face protection.

FIRST AID:

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a POISON CENTER or doctor/physician. Collect spillage.

SAFETY DATA SHEET:

For additional information, refer to SDS available from your supplier

CONDITIONS OF SALE - LIMITED WARRANTY AND LIMITATIONS OF LIABILITY AND REMEDIES

Read the Conditions of Sale - Warranty and Limitations of Liability and Remedies before using this product. If the terms are not acceptable, return the product, unopened, and the full purchase price will be refunded. This label is believed to be reliable and should be followed carefully. Injury to the crop to which the product is applied may result from the occurrence of extraordinary or unusual weather conditions or the failure to follow the label directions or good application practices, all of which are beyond the control of Campbells Fertilisers Australasia Pty Ltd (the "Company") or seller. In addition, failure to follow label directions may cause injury to crops, animals, man or the environment. The Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purpose referred to subject to the factors noted above which are beyond the control of the Company. The Company makes no other warranties or representations of any kind, express or implied, concerning the product, including no implied warranty of merchantability or fitness for any particular purpose, and no such warranty shall be implied by law. The exclusive remedy against the Company for any cause of action relating to the handling or use of this product shall be limited to, at Campbells Fertilisers Australasia's election, one of the following: 1. Refund of the purchase price paid by buyer or user for product bought, or 2. Replacement of the product used. To the extent allowed by law, the Company shall not be liable and any and all claims against the Company are waived for special, indirect, incidental, or consequential damages or expense of any nature, including, but not limited to, loss of profits or income. The Company and the seller offer this product and the buyer and user accept it, subject to the foregoing conditions of sale and limitation of warranty, liability and remedies.

MANUFACTURED FOR Campbells Fertilisers Australasia Pty Ltd

CAUTION:

BEFORE USING THIS PRODUCT READ ALL PRECAUTIONS, DIRECTIONS FOR USE, CONDITIONS OF SALE-LIMITED WARRANTY AND LIMITATIONS OF LIABILITY AND REMEDIES. Avoid contact with formulated product. Do not take internally. Avoid contact with or inhalation of the spray mist. Do not apply this product in such a manner as to directly expose workers or other persons. If product is being mixed with pesticides and/or spray adjuvants, follow all precautionary statements on the accompanying product(s) labelling. Avoid contact with eyes and prolonged contact with skin to avoid irritation. Wear safety glasses and chemical resistant gloves. Not for human or animal consumption.

GENERAL INFORMATION This product is recommended for use as a supplement to a regular fertilizer program only.

Causes skin irritation. Causes serious eye irritation. May be harmful if swallowed, in contact with skin or if inhaled. Do not get in eyes, on skin, or on clothing. Wash contacted areas thoroughly after handling. Wear protective gloves, protective clothing and eye or face protection. Take off contaminated clothing and wash before reuse. IF SWALLOWED: Rinse mouth. Do NOT induce vomiting. IF ON SKIN: Wash with plenty of soap and water. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If skin irritation occurs: Get medical advice. If eye irritation persists: Get medical advice or attention. In case of fire: Use carbon dioxide, dry chemical, foam, water fog, to extinguish. Protect from sunlight. Store in a dry place. Store in a closed container. Store in a well-ventilated place. Keep cool. If they can not be recycled, dispose of contents to an approved waste disposal plant and containers to landfill (see Section 13 of this SDS).

**IN A TRANSPORT EMERGENCY
DIAL 000
POLICE OR FIRE BRIGADE**

DISTRIBUTED BY:

Campbells Fertiliser Australasia Pty Ltd

18 Raymond Road Laverton North Victoria 3026 Australia

Phone: +61 3 9931 2211 Fax: +61 3 9931 2201 www.campbellsfert.com.au



AUTHORISED DISTRIBUTOR

Is a registered trademark of ®CoRoN is a registered trademark of Helena Holding Company